

A row of microdroppers, likely from a liquid handling robot, is shown dispensing a blue liquid into a 96-well plate. The droppers are in the foreground, and the wells of the plate are visible in the background, creating a sense of depth and precision in a laboratory setting.


Building Synthetic Circuits.

Nicola J Patron | Plant Sciences, University of Cambridge
[http:// patronlab.org](http://patronlab.org)

Learning Goals


To understand:

- How to decide when and what to synthesise
- What are standardised DNA parts
- How to assemble multiple DNA parts in a single reaction
- How standardisation aids construct design and assembly




Synthesis vs. Assembly

Synthesis – oligos, blocks & genes





Abstraction



Figures: Patron (2020) New Phytologist

Standardisation

Biological Standards:
The BioBrick (BBF RFC10)
2004



Registry of Standard
Biological Parts

Standardisation

cannot , do
I not understand
what
create I

Richard Feynman

Standardisation

Biological Standards: Common Genetic Syntax (BBF RFC106) 2015





IMAGE: iGEM


Overlap-dependent Parallel Assembly

Isothermal (aka Gibson) Assembly


Production of overlapping double-stranded DNA fragments



Overlapping linear fragments are combined with a three-enzyme cocktail:



T5 exonuclease performs a 5'-3' resection of all components allowing complementary overhangs to self-anneal.
A proof-reading **DNA polymerase** & **Taq-ligase** fill in and join gaps



Overlap-dependent Parallel Assembly

16,520 bp mouse mitochondrial genome from 600 fragments





Figure: Gibson et al (2010) Nature Methods 7, 901–903

Ligase-dependent Parallel Assembly

Type IIS (aka Golden Gate) Assembly




Ligase-dependent Parallel Assembly

Type II Restriction Endonucleases

Palindromic, cleave inside of their recognition sequence:

EcoRI



G AATTC



4 base pair 5' overhang

ApaI


GGGCC C



4 base pair 3' overhang

ScaI

AGT ACT




Ligase-dependent Parallel Assembly

Type II^S Restriction Endonucleases

Non-palindromic, cleave **outside** of their recognition sequence:

Bpil GAAGACNN□NNNN




4 base pair 5' overhang

Ligase-dependent Parallel Assembly

Type II^S Restriction Endonucleases

Non-palindromic, cleave **outside** of their recognition sequence:



No recognition-sequence
in released fragment




Ligase-dependent Parallel Assembly

Multiple fragments can be assembled in a desired order

Acceptor with DIVERGENT recognition sequences




Parts with CONVERGENT recognition sequences




4 (ii) Ligase-dependent Parallel Assembly

Multiple fragments can be assembled in a desired order



Ligase-dependent Parallel Assembly

Multiple fragments can be assembled in a desired order
Recognition sequence is not present in the ligated product




Ligase-dependent Parallel Assembly


One-step Type IIS Digestion-Ligation (Dig-Lig)

Since the recognition sequence is never present in the ligated product:

- The digestion and ligation are performed simultaneously
- The reaction can be repetitively cycled for maximum efficiency



Standardisation + Abstraction + Automation = Scalability



- Increases throughput - many experiments performed simultaneously
- Reduces reaction volumes
- Increases reproducibility by eliminating many human errors

END OF PART

Building Synthetic Circuits

- How to decide when and what to synthesise
- What are standardised DNA parts
- How to assemble multiple DNA parts in a single reaction
- How standardisation aids construct design and assembly